

**ENVIRONMENTAL-RISK (E-RISK) LONGITUDINAL TWIN STUDY
CONCEPT PAPER FORM**

Proposing Authors: Timothy Matthews / Line Jee Hartmann Rasmussen

Author's affiliation, phone, and e-mail address:

Timothy Matthews:

Social, Genetic, and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London, United Kingdom

timothy.matthews@kcl.ac.uk

Line J. H. Rasmussen:

Department of Psychology and Neuroscience, Duke University, Durham, NC

Clinical Research Centre, Copenhagen University Hospital Hvidovre, Hvidovre, Denmark

line.jee.hartmann.rasmussen@duke.edu

Sponsoring Investigator (if the proposing author is a student, a post-doc or a colleague):

Louise Arsenaault

Terrie E. Moffitt

Avshalom Caspi

Proposed co-authors:

Andrea Danese

Jesper Eugen-Olsen

Daisy Fancourt

Karen Sugden

Benjamin Williams

Antony Ambler

Richie Poulton

Kasper Iversen

Martin Schultz

Provisional Paper Title:

Social isolation, loneliness, and inflammation

Date:

9 October, 2019

AMENDMENT: 5 March, 2020

Due to availability of new data, we have made an amendment to the concept paper, please see changes made in red.

Objective of the study and its significance:

The importance of social relationships for health and longevity is supported by a large and growing body of evidence (Umberson & Montez, 2010). According to a recent meta-analysis, social isolation, loneliness and living alone are each associated with a 25-30% increase in risk for mortality (Holt-Lunstad et al, 2015). Numerous mechanisms for this association have been proposed, including maladaptive health-related behaviours, increased blood pressure and impaired sleep quality.

Another pathway through which social relationships could influence health is inflammation. According to a hypothesis based on evolutionary theory (Cacioppo et al, 2006), loneliness is an adaptive response to social disconnection that prepares individuals to face an unsafe environment without protection from kith

and kin. Based on this premise, it is hypothesised that loneliness is accompanied by changes in immune functioning that would bolster an individual's ability to fight infection in event of wounding. Consistent with this, individuals high in feelings of loneliness exhibit a pattern of pro-inflammatory changes in gene expression (Cole et al, 2007).

A limitation of the existing literature on social connection and inflammation is that studies have often conflated social isolation and loneliness, or only examined one of the two. These are frequently co-occurring but conceptually distinct phenomena, and it is not clear which is the 'active ingredient'. Those studies that have examined both in parallel (e.g. Shankar et al, 2011) suggest that objective social isolation, rather than loneliness, is the more relevant risk factor for this particular outcome. However, this research has been cross-sectional in nature and cannot address the possibility that individuals suffering from inflammatory disease may, as a result, be restricted from engaging in social activities. Longitudinal studies have found that childhood social isolation predicts increased inflammation in adults (Danese et al, 2009; Lacey et al, 2014), but there is limited longitudinal research on loneliness to compare this to. A further issue still is that loneliness is highly correlated with depression (Matthews et al, 2016), and studies that fail to take their co-occurrence into account may yield associations that are confounded.

There are several ways to assess systemic inflammation, each with advantages and disadvantages in their cost, utility, and predictive validity. We have recently identified one such marker of systemic inflammation, suPAR (*soluble urokinase plasminogen activator receptor*), as a more effective measure of chronic stress than other, more traditional measures of systemic inflammation used in clinical or research settings, i.e., high-sensitivity C-reactive protein (hsCRP) and interleukin-6 (IL-6). Specifically, we found increased suPAR levels in individuals who had been exposed to early life risk factors, such as adverse childhood experiences, or specific victimization experiences (Rasmussen et al., 2019a; 2019b). In contrast to CRP and IL-6, both of which are acute-phase reactants and markers of acute inflammation and infections (Rhodes et al., 2011; Hunter & Jones, 2015), suPAR appears to be less affected by acute conditions (Lyngbæk et al., 2012) and therefore potentially a better measure of systemic chronic inflammation. The plasma level of suPAR is thought to reflect a person's overall level of immune activity, and elevated suPAR is observed in a wide range of diseases and pathological conditions (Hayek et al., 2015; Persson et al., 2014; Schaefer et al., 2017; Theilade et al., 2015). Research indicates that suPAR is associated with clinical outcomes independent of CRP and IL-6 (Rasmussen et al., 2016; Botha et al., 2015).

Here, we wish to use data from **the Danish clinical sample TRIAGE I and the longitudinal population-representative E-Risk and Dunedin studies** to investigate the associations between social isolation and loneliness with markers of systemic inflammation, including the acute inflammation markers CRP and IL-6 and the chronic inflammation marker suPAR.

Statistical analyses:

Aim 1 (primary analysis): Investigate the association of childhood social isolation and adolescent/adulthood loneliness with suPAR, hsCRP, and IL-6 levels at age 18 in the E-Risk Study and at age 38 in the Dunedin Study, in separate models of social isolation or loneliness, and in models including both social isolation and loneliness. **Moreover, investigate the association of childhood social isolation and adulthood loneliness at age 38 with suPAR, hsCRP, and IL-6 levels at age 45 in the Dunedin Study.**

Aim 2 (secondary analysis): Investigate the association of depression with suPAR, hsCRP, and IL-6 levels at age 18 in the E-Risk Study and at age 38 in the Dunedin Study. Test whether the associations found in Aim 1 are robust to controls for depression.

Aim 3: Investigate the association of "living alone" with suPAR, CRP, and IL-6 in the Danish TRIAGE I study.

Associations will be tested using multivariable linear regression analyses. All models will include a variety of relevant demographic, childhood, and health behaviour covariates, such as sex, body mass index, and smoking. Further adjustments will be made for childhood SES. A p value < 0.05 is a priori designated as statistically significant.

Variables Needed at Which Ages (names and labels):		
	E-Risk:	Dunedin:
General:	familyid Unique family identifier atwinid Twin A ID (ex chkdg) btwinid Twin B ID (ex chkdg) rorderp5 Random Twin Order risks Sample Groups cohort Cohort sampsex Sex of Twins: In sample zygosity Zygosity	Sex
Childhood:	Age 5: sisoe5 Social isolation scale (mum & teacher average) - P5 – Elder sisoy5 Social isolation scale (mum & teacher average) - P5 – Younger Age 7: sisoe7 Social isolation scale (mum & teacher average) – P7 – Elder sisoy7 Social isolation scale (mum & teacher average) – P7 - Younger Age 10: sisoe10 Social isolation scale (mum & teacher average) – P10 - Elder sisoy10 Social isolation scale (mum & teacher average) – P10 - Younger Age 12: sisoe12 Social isolation scale (mum & teacher average) – P12 – Elder sisoy12 Social isolation scale (mum & teacher average) – P12 – Younger lonelye12 Loneliness – P12 – Elder lonelyy12 Loneliness – P12 – Younger cdie12 Depression scale – CDI – Elder cdiy12 Depression scale – CDI - Younger	Social isolation (Zisolkid)
Age 18:	socisoe18 Social Isolation scale - P18 – Elder socisoy18 Social Isolation scale - P18 – Younger lonelye18 Loneliness scale – P18 – Elder lonelyy18 Loneliness scale – P18 – Younger mdesxe18 MDE – symptom scale – P18 – Elder mdesxy18 MDE – symptom scale – P18 – Younger dxmdee18 Major depressive episode, dsm4 – P18 – Elder dxmdey18 Major depressive episode, dsm4 – P18 - Younger suPAR (suPARngml_E, suPARngml_Y) CRP (CRPmgLE18, CRPmgLY18) IL-6 (IL6pgmLE18, IL6pgmLY18)	

	Smoking: - smkcure18, smkcury18 - smkcnume18, smkcnumy18 - smkpyre18, smkpyry18 Body mass index (bmie18, bmiy18)	
Age 38:		suPAR (suPAR38ngml) CRP (CRP38np, LNCRP38np) IL-6 () Smoking (Smoker38, PackYrLifTm38, CigsDay38) Body mass index (bmi38np) Current illness (SickToday38, sickat38) Loneliness (Lonely38, LonelyUCLA38)
Age 45:		suPAR at age 45 CRP at age 45 IL-6 at age 45 Current smoking Body mass index Current illness Loneliness (LonelyUCLA45) Depression (dxmde45)

Data from TRIAGE I:

- Sex
- Age
- BMI
- CRP
- IL-6
- suPAR
- Living alone (y/n)
- Living in nursing home (y/n)
- Current smoking (yes, former, never, unknown)
- Mortality
- Available diagnoses

References cited:

Botha, S., Fourie, C.M., Schutte, R., Eugen-Olsen, J., Pretorius, R., & Schutte, A.E. (2015) Soluble urokinase plasminogen activator receptor as a prognostic marker of all-cause and cardiovascular mortality in a black population. *Int J Cardiol*, 184:631-636.

Cacioppo, J. T., Hawkey, L. C., Ernst, J. M., Burleson, M., Berntson, G. G., Nouriani, B., & Spiegel, D. (2006). Loneliness within a nomological net: An evolutionary perspective. *J Res Pers*, 40(6), 1054-1085.

Cole, S. W., Hawkey, L. C., Arevalo, J. M., Sung, C. Y., Rose, R. M., & Cacioppo, J. T. (2007). Social regulation of gene expression in human leukocytes. *Genome Biol*, 8(9), R189.

Hayek, S.S., Sever, S., Ko, Y.A., et al. (2015) Soluble Urokinase Receptor and Chronic Kidney Disease. *N Engl J Med*, 373(20):1916-1925.

Holt-Lunstad, J., Smith, T. B., Baker, M., & Stephenson, D. (2015). Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspect Psychol Sci*, 10(2), 227-237.

Hunter, C.A., & Jones, S.A. (2015) IL-6 as a keystone cytokine in health and disease. *Nat Immunol*, 16(5):448-457.

Lacey, R. E., Kumari, M., & Bartley, M. (2014). Social isolation in childhood and adult inflammation: evidence from the National Child Development Study. *Psychoneuroendocrinology*, 50, 85-94.

- Lyngbæk, S., Marott, J.L., Møller, D.V., et al. (2012) Usefulness of soluble urokinase plasminogen activator receptor to predict repeat myocardial infarction and mortality in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous intervention. *Am J Cardiol*, 110(12):1756-1763.
- Matthews, T., Danese, A., Wertz, J., Odgers, C., Ambler, A., Moffitt, T. E., & Arseneault L. (2016). Social isolation, loneliness and depression in young adulthood: A behavioural genetic analysis. *Social Psychiatry and Psychiatric Epidemiology*, 51(3), 339-348.
- Persson, M., Östling, G., Smith, G., et al. (2014) Soluble Urokinase Plasminogen Activator Receptor: A Risk Factor for Carotid Plaque, Stroke, and Coronary Artery Disease. *Stroke*, 45(1):18-23.
- Rasmussen, L.J.H., Ladelund, S., Haupt, T.H., et al. (2016) Soluble urokinase plasminogen activator receptor (suPAR) in acute care: A strong marker of disease presence and severity, readmission and mortality. A retrospective cohort study. *Emerg Med J*, 33(11):769-775.
- Rasmussen, L.J.H., Moffitt, T.E., Eugen-Olsen, J., Belsky, D.W., Danese, A., Harrington, H.L., Houts, R.M., Poulton, R., Sugden, K., Williams, B.S., & Caspi, A. (2019a). Cumulative childhood risk is associated with a new measure of chronic inflammation in adulthood. *J Child Psychol Psychiatr*, 60(2), 199-208.
- Rasmussen, L.J.H., Moffitt, T.E., Arseneault, L., Danese, A., Eugen-Olsen, J., Fisher, H.L., Harrington, H.L., Houts, R., Matthews, T., Sugden, K., Williams, B.S., & Caspi, A. (2019b) Improving the measurement of stress-related inflammatory burden in young people: A longitudinal cohort study. In Press.
- Rhodes, B., Fürnrohr, B.G., Vyse, T.J. (2011) C-reactive protein in rheumatology: Biology and genetics. *Nat Rev Rheumatol*, 7(5):282-289.
- Schaefer, F., Trachtman, H., Wühl, E., et al. (2017) Association of serum soluble urokinase receptor levels with progression of kidney disease in children. *JAMA Pediatr*, 171(11):e172914.
- Shankar, A., McMunn, A., Banks, J., & Steptoe, A. (2011). Loneliness, social isolation, and behavioral and biological health indicators in older adults. *Health Psychol*, 30(4), 37-385.
- Theilade, S., Lyngbaek, S., Hansen, T.W., et al. (2015) Soluble urokinase plasminogen activator receptor levels are elevated and associated with complications in patients with type 1 diabetes. *J Intern Med*, 277(3):362-371.
- Umberson, D. & Montez, J. K. (2010). Social relationships and health: a flashpoint for health policy. *J Health Soc Behav*, 51(Suppl), S54-266.

Data Security Agreement

Provisional Paper Title	Social isolation, loneliness, and inflammation
Proposing Author	Timothy Matthews / Line Jee Hartmann Rasmussen
Today's Date	17 September 2019

Please keep one copy for your records

(Please initial your agreement)

TM/LR I am familiar with the King's College London research ethics guidelines (<https://www.kcl.ac.uk/innovation/research/support/ethics/about/index.aspx>) and the MRC good research practice guidelines (<https://www.mrc.ac.uk/research/policies-and-guidance-for-researchers/good-research-practice/>).

TM/LR My project has ethical approval from my institution.

TM/LR I am familiar with the EU General Data Protection Regulation (<https://mrc.ukri.org/documents/pdf/gdpr-guidance-note-3-consent-in-research-and-confidentiality/>), and will use the data in a manner compliant with its requirements.

TM/LR My computer is (a) encrypted at the hard drive level, (b) password-protected, (c) configured to lock after 15 minutes of inactivity, AND (d) has an antivirus client which is updated regularly.

TM/LR I will treat all data as "restricted" and store in a secure fashion.

TM/LR I will not share the data with anyone, including students or other collaborators not specifically listed on this concept paper.

TM/LR I will not merge data from different files or sources, except where approval has been given by the PI.

TM/LR I will not post data online or submit the data file to a journal for them to post. Some journals are now requesting the data file as part of the manuscript submission process. The E-Risk Study cannot be shared because the Study Members have not given informed consent for unrestricted open access. Speak to the study PI for strategies for dealing with data sharing requests from Journals.

TM/LR Before submitting my paper to a journal, I will submit my draft manuscript and scripts for data checking, and my draft manuscript for co-author mock review, allowing three weeks.

TM/LR I will submit analysis scripts and new variable documentation to project data manager after the manuscript gets accepted for publication.

TM/LR I will delete the data after the project is complete.

TM/LR **For projects using location data:** I will ensure geographical location information, including postcodes or geographical coordinates for the E-Risk study member's homes or schools, is never combined or stored with any other E-Risk data (family or twin-level data)

TM/LR **For projects using genomic data:** I will only use the SNP and/or 450K data in conjunction with the phenotypes that have been approved for use in this project at the concept paper stage.

Signature:



CONCEPT PAPER RESPONSE FORM

A. To be completed by the proposing author

Proposing Author: Timothy Matthews / Line Jee Hartmann Rasmussen

X I have read the E-Risk data-sharing policy guidelines and agree to follow them

Provisional Paper Title: Social isolation, loneliness, and inflammation

Potential co-authors: Louise Arseneault, Terrie E. Moffitt, Avshalom Caspi, Andrea Danese, Jesper Eugen-Olsen, Daisy Fancourt, Kasper Iversen, Martin Schultz, Karen Sugden, Benjamin Williams, Antony Ambler, Richie Poulton

Potential Journals: TBC

Intended Submission Date (month/year): Summer 2020

Please keep one copy for your records and return one to Louise (louise.arseneault@kcl.ac.uk)

B. To be completed by potential co-authors:

Approved Not Approved Let's discuss, I have concerns

Comments:

Please check your contribution(s) for authorship:

- Conceptualizing and designing the longitudinal study
- Conceptualizing and collecting one or more variables
- Data collection
- Conceptualizing and designing this specific paper project
- Statistical analyses
- Writing
- Reviewing manuscript drafts
- Final approval before submission for publication
- Acknowledgment only, I will not be a co-author

Signature:

